

Further Maths  
A-Level Starter  
Activity



**Topic: The Method of Differences (1)**  
Chapter Reference: Core Pure 2, Chapter 2

**10  
minutes**

1. Given that  $\frac{r+1}{r+2} - \frac{r}{r+1} = \frac{1}{(r+1)(r+2)}$ , find an expression, in terms of  $n$ , for (4)

$$\frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \dots + \frac{1}{(n+1)(n+2)}$$

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

2. Given that  $\frac{1}{r} - \frac{1}{r+2} = \frac{2}{r(r+2)}$ , find an expression, in terms of  $n$ , for (5)

$$\frac{2}{1 \times 3} + \frac{2}{2 \times 4} + \dots + \frac{2}{n(n+2)}$$

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



## Solutions

1. Either

$\frac{2}{3} - \frac{1}{2} + \frac{3}{4} - \frac{2}{3} \dots \frac{n+1}{n+2} - \frac{n}{n+1}$ Express terms as differences	<b>M1</b>
At least first two and last term correct	<b>A1</b>
Show or imply that pairs of terms cancel	<b>M1</b>
$\frac{n+1}{n+2} - \frac{1}{2}$ Obtain correct answer in any form	<b>A1</b>

Or

State that $\sum_{r=1}^n u_r = f(n+1) - f(1)$	<b>M2</b>
Each term correct	<b>A1A1</b>

2.

Express terms as differences	<b>M1</b>
Express 1 <sup>st</sup> 3 (or last 3) terms so that cancelling occurs	<b>M1</b>
Obtain $1 + \frac{1}{2}$	<b>A1</b>
Obtain $-\frac{1}{n+2}, -\frac{1}{n+1}$	<b>A1</b>
$\frac{3}{2} - \frac{1}{n+1} - \frac{1}{n+2}$ Obtain correct answer in any form	<b>A1</b>





## Solutions

1.

a	$= (2x + 1)e^{2x}$	<b>A2</b>
b	$= (4x + 4)e^{2x}$	<b>A2</b>
c	$= (8x + 12)e^{2x}$	<b>A2</b>
d	$= (16x + 32)e^{2x}$	<b>A2</b>
e	$= (32x + 80)e^{2x}$	<b>A2</b>





## Solutions

1a.

Attempt quotient/product on bracket	<b>M1</b>
Get $-3/(2+x)^2$ May be implied	<b>A1</b>
Use Formulae Booklet or derive from $\tanh y = (1-x)/(2+x)$ Attempt $\tanh^{-1}$ part in terms of $x$	<b>M1</b>
Get $\frac{-3}{(2+x)^2} \cdot \frac{1}{1 - ((1-x)/(2+x))^2}$	<b>A1</b>
Clearly tidy to A.G.	<b>A1</b>
Get $f''(x) = 2/(1+2x)^2$	<b>B1</b>

OR

Use reasonable ln definition	<b>M1</b>
Get $y = \frac{1}{2} \ln((1-k)/(1+k))$ for $k = (1-x)/(1+2x)$ .	<b>A1</b>
Tidy to $y = \frac{1}{2} \ln(3/(1+2x))$	<b>A1</b>
Attempt chain rule	<b>M1</b>
Clearly tidy to A.G.	<b>A1</b>
Get $f''(x)$	<b>B1</b>

1b.

Attempt $f(0)$ , $f'(0)$ and $f''(0)$	<b>M1</b>
Get $\tanh^{-1} \frac{1}{2}$ , $-1$ and $2$	<b>A1</b>
Replace $\tanh^{-1} \frac{1}{2} = \frac{1}{2} \ln 3 (= \ln \sqrt{3})$	<b>B1</b>
Get $\ln \sqrt{3} - x + x^2$	<b>A1</b>





## Solutions

1a.

Substitution of $2x$	<b>A1</b>
Use of standard Maclaurin expansion formula	<b>M1</b>
$2x - 8x^2 + \frac{26}{3}x^3 - 12x^4 + \dots$	<b>A1</b>

1b.

Substitution of $2x$	<b>A1</b>
Use of standard Maclaurin expansion formula	<b>M1</b>
$x + 2x^2 + \frac{11}{6}x^3 + x^4 + \dots$	<b>A1</b>

1c.

Substitution of $x^2$	<b>A1</b>
Use of standard binomial expansion formula	<b>M1</b>
$1 - x + x^2 - \frac{2}{3}x^3 + \frac{1}{6}x^4 + \dots$	<b>A1</b>







## Solutions

1a.

Reasonable attempt at chain rule	<b>M1</b>
Reasonable attempt at product/quotient rule	<b>M1</b>
Correctly get $f'(0) = 1$	<b>A1</b>
Correctly get $f''(0) = 1$	<b>A1</b>

1b.

Reasonable attempt at Maclaurin with their values	<b>M1</b>
Get $1 + x + \frac{1}{2}x^2$	<b>A1</b>

