

**Further Maths
A-Level Starter
Activity**



Topic: Hooke's Law (1)

Chapter Reference: Further Mechanics 1, Chapter 3

**10
minutes**

1. The force required to stretch Hooke's Law spring varies from 0 N to 65 N as we stretch the spring by moving one end 6.3 cm from its unstressed position. Find the force constant of the spring. Answer in units of N/m. (3)

2. A string with modulus (of elasticity) 10 N has a natural length of 2m. What is the tension in the string when its length is 5m. (2)

3. A 75 kg person stands on a compression spring with spring constant 5000 N/m and nominal length 0.25m. What is the total length of the loaded spring. (5)

Don't know where to start? Draw a daigram



Solutions

1.

$6.3 \text{ cm} = 0.063 \text{ m}$ $65 = k \times 0.063$	M1
$k = \frac{65}{0.063}$	M1
$= 1032 \text{ N/m}$	A1

2.

$T = \frac{10 \times 3}{2}$	M1
$T = 15 \text{ N}$	A1

3.


$x = \frac{F}{k} = \frac{mg}{k}$	M1
$x = \frac{75 \times 9.81}{5000}$	M1
$x = 0.15 \text{ m}$	A1
Therefore, length of loaded spring = Original length - x	M1
$= 0.1 \text{ m}$	A1



Solutions

1.


Strings are light \rightarrow ignore masses and assume tensions are equal.



Resolving vertically	M1
$T = 3g$	A1
For S_1 : $T = 3g = \frac{\lambda x}{l} = \frac{75x_1}{0.5}$	M1
$x_1 = \frac{g}{50}$	A1
For S_2 : $T = 3g = \frac{\lambda x}{l} = \frac{120x_2}{0.4}$	M1
$x_2 = \frac{g}{100}$	A1
$x_1 + x_2 = \frac{3g}{100} = 0.295$	M1
Distance of 3kg mass below O, is $0.5 + 0.4 + 0.294$	M1
$= 1.194 = 1.2\text{m}$	A1

Solutions

1a.



Resolving vertically

M1

$$2T \cos \theta = W$$

A1

$$\text{Hooke's Law: } T = \frac{80 \times 3.5}{4}$$

M1

Therefore $W = 84 \text{ N}$

A1

1b.

$$\text{Elastic Potential Energy} = 2 \times \frac{80 \times 3.5^2}{2 \times 4}$$

M1

$$= 245$$

A1

Solutions

1a.

$\text{EPE} = \frac{\lambda \times 0.5^2}{1.2}$	B1
$\text{GPE lost} = \text{EPE gained}$	M1
$0.8 \times 9.8 \times 1.1 = \frac{\lambda \times 0.5^2}{1.2}$	A1
$\lambda = 41.4 \text{ N or } 41 \text{ N}$	A1



Solutions

1a.

Elastic Potential Energy = $\frac{1}{2} \frac{3.6mg}{a} x^2 = \frac{1}{2} \frac{3.6mg}{a} \left(\frac{a}{3}\right)^2$	M1 A1
= 0.2 mga	A1

1b.

Friction = μmg	M1
Therefore, work done by friction = $\mu mg \left(\frac{4a}{3}\right)$	M1
Work-energy = $\frac{1}{2} m 2ga = \mu mgd + 0.2 mga$	M1 M1
Solving: $\mu = 0.6$	A1

