



1. The quadratic equation

$$x^2 + px + q = 0$$

in which the coefficients  $p$  and  $q$  are real, has a complex root  $\sqrt{5} - i$ .

a. Write down the other root of the equation

(1)

---

---

---

---

---

b. Find the sum and product of the two roots of the equation.

(3)

---

---

---

---

---

c. Hence state the values of  $p$  and  $q$ .

(2)

---

---

---

---

---



## Solutions

1a.

$\sqrt{5} + i$	<b>B1</b>
----------------	-----------

1b.

Sum of roots is $2\sqrt{5}$ Product is 6	<b>B1</b> <b>M1A1</b>
---	--------------------------

1c.

$p = -2\sqrt{5}, q = 6$	<b>B1</b> <b>B1</b>
-------------------------	------------------------



**Further Maths  
A-Level Starter  
Activity**



**Topic: Roots of Polynomials (2)**

Chapter Reference: Core Pure 1, Chapter 4

**10  
minutes**

1. The cubic equation  $3x^3 - 9x^2 + 6x + 2 = 0$  has roots  $\alpha$ ,  $\beta$  and  $\gamma$ .

Write down the values of  $\alpha + \beta + \gamma$ ,  $\alpha\beta + \beta\gamma + \gamma\alpha$  and  $\alpha\beta\gamma$ .

**(3)**

---

---

---

---

---

---

---

2. The cubic equation  $3z^3 + pz^2 + 17z + q = 0$ , where  $p$  and  $q$  are real, has a root  $\alpha = 1 + 2i$ .

a. Write down the value of another non-real root,  $\beta$ , of this equation.

**(1)**

---

---

---

b. Find the value of the third root,  $\gamma$ , of this equation.

**(3)**

---

---

---

---

---

---

---

c. Find the values of  $p$  and  $q$ .

**(3)**

---

---

---

---

---

---

---



## Solutions

1.

$\alpha + \beta + \gamma = 3$ $\alpha\beta + \beta\gamma + \gamma\alpha = 2$ $\alpha\beta\gamma = \frac{2}{3}$	<b>B1</b> <b>B1</b> <b>B1</b>
--	-------------------------------------

2a.

$1 - 2i$	<b>B1</b>
----------	-----------

2b.

$\sum \alpha\beta = \frac{17}{3}$ $\alpha\beta + \beta\gamma + \gamma\alpha = \frac{17}{3}$ $\Rightarrow \gamma = \frac{1}{3}$	<b>B1</b> <b>M1</b> <b>A1</b>
--	-------------------------------------

2c.

$\alpha + \beta + \gamma = \frac{-p}{3}, \quad \alpha\beta\gamma = \frac{-q}{3}$ $p = -7$ $q = -5$	<b>M1</b> <b>A1</b> <b>A1</b>
--	-------------------------------------





1. The equation,

$$x^4 - 6x^3 - 73x^2 + kx + m = 0$$

has two positive roots  $\alpha, \beta$  and two negative roots  $\gamma, \delta$ . It is given that  $\alpha\beta = \gamma\delta = 4$ .

a. Find the values of the constant  $k$  and  $m$ .

(5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

b. Show that  $(\alpha + \beta)(\gamma + \delta) = -81$

(4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



## Solutions

1a.

$\alpha + \beta + \gamma + \delta = 6$ $k = -(\beta\gamma\delta + \alpha\gamma\delta + \alpha\beta\delta + \alpha\beta\gamma)$ $= -4(\beta + \alpha + \delta + \gamma)$ $= -24$ $m = \alpha\beta\gamma\delta = 16$	<b>B1</b> <b>M1</b> <b>M1</b> <b>A1</b> <b>B1</b>
--	---

1b.

$\sum\alpha\beta = -73$ $(\alpha + \beta)(\gamma + \delta) = \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta$ $\sum\alpha\beta - \alpha\beta - \gamma\delta$ $= -73 - 4 - 4$ $= -81$	<b>B1</b> <b>M1</b> <b>A1</b> <b>A1</b>
--	--





1. Show that  $(\alpha\beta + \beta\gamma + \gamma\alpha)^2 \equiv \alpha^2\beta^2 + \beta^2\gamma^2 + \gamma^2\alpha^2 + 2\alpha\beta\gamma(\alpha + \beta + \gamma)$ . (3)

---

---

---

---

---

---

---

---

---

---

2. It is given that  $\alpha$ ,  $\beta$  and  $\gamma$  are the roots of the cubic equation  $x^3 + px^2 - 4x + 3 = 0$ , where  $p$  is a constant.  
Find the value of  $\frac{1}{\alpha^2} + \frac{1}{\beta^2} + \frac{1}{\gamma^2}$  in terms of  $p$ . (5)

---

---

---

---

---

---

---

---

---

---



## Solutions

1.

Attempt at complete expansion	<b>M1</b>
Obtain correct unsimplified answer	<b>A1</b>
Obtain <b>given</b> answer correctly	<b>A1</b>

2.

$\Sigma\alpha = -p, \Sigma\alpha\beta = -4, \alpha\beta\gamma = -3$	<b>B1</b>
	<b>M1</b>
$\frac{16 - 6p}{9}$	<b>A1</b>
	<b>M1</b>
	<b>A1</b>







1. The cubic equation  $3x^3 - 9x^2 + 6x + 2 = 0$  has root  $\alpha$ ,  $\beta$  and  $\gamma$ .

a. Write down the values of  $\alpha + \beta + \gamma$  and  $\alpha\beta + \beta\gamma + \gamma\alpha$ . (2)

---

---

---

---

---

b. Find the value of  $\alpha^2 + \beta^2 + \gamma^2$ . (2)

---

---

---

---

---

c. Use the substitution  $x = \frac{1}{u}$  to find a cubic equation in  $u$  with integer coefficients. (2)

---

---

---

---

---

d. Use your answer to part (c) to find the value of  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$ . (2)

---

---

---

---

---



## Solutions

1a.

$\alpha + \beta + \gamma = 3, \alpha\beta + \beta\gamma + \gamma\alpha = 2$	<b>B1</b> <b>B1</b>
---	------------------------

1b.

$\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ $= 9 - 4 = 5$	<b>M1</b> <b>A1</b>
--	------------------------

1c.

$\frac{3}{u^3} - \frac{9}{u^2} + \frac{6}{u} + 2 = 0$ $2u^3 + 6u^2 - 9u + 3 = 0$	<b>M1</b> <b>M1</b>
---	------------------------

1d.

$\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = -3$	<b>M1</b> <b>A1</b>
--	------------------------

