



# Practice Exam Paper H

Time: 2 Hours



1. The line  $L$  has equation  $y = 5 - 2x$

a. Show that the point  $P(3, -1)$  lies on  $L$  (1)

b. Find an equation of the line perpendicular to  $L$ , which passes through  $P$ . Give your answer in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers. (4)

**(Total Marks: 5)**

2. Write  $\sqrt{45}$  in the form  $a\sqrt{5}$ , where  $a$  is an integer. (1)

b. Express  $\frac{2(3+\sqrt{5})}{(3-\sqrt{5})}$  in the form  $b + c\sqrt{5}$ , where  $b$  and  $c$  are integers. (5)

**(Total Marks: 6)**

3. Given that  $f(x) = \frac{1}{x}$ ,  $x \neq 0$

a. Sketch the graph of  $y = f(x) + 3$  and state the equations of the asymptotes. (4)

b. Find the coordinates of the point where  $y = f(x) + 3$  crosses a coordinate axis. (2)

**(Total Marks: 6)**

4. The equation  $2x^2 - 3x - (k + 1) = 0$ , where  $k$  is a constant, has no real roots.

Find the set of possible values of  $k$  (4)

**(Total Marks: 4)**

5. The curve  $C$  has equation  $y = f(x)$ ,  $x > 0$ , and  $f'(x) = 4x - 6\sqrt{x} + \frac{8}{x^2}$

Given that the point  $P(4, 1)$  lies on  $C$ ,

a. Find  $f(x)$  and simplify your answer. (6)

b. Find an equation of the normal to  $C$  at the point  $P(4, 1)$ . (4)

**(Total Marks: 10)**

6.  $f(x) = x^2 + 4kx + (3 + 11k)$ , where  $k$  is a constant.

a. Express  $f(x)$  in the form  $(x + p)^2 + q$ , where  $p$  and  $q$  are constants to be found in terms of  $k$ . (3)

Given that the equation  $f(x) = 0$  has no real roots

b. Find the set of possible values of  $k$ . (4)

Given that  $k = 1$

c. Sketch the graph of  $y = f(x)$ , showing the coordinates of any point at which the graph crosses a coordinate axis. (3)

**(Total Marks: 13)**

7. The points  $A$  and  $B$  have coordinates  $(5, -1)$  and  $(13, 11)$  respectively.

a. Find the coordinates of the mid-point of  $AB$ . (2)

Given that  $AB$  is a diameter of the circle  $C$ ,

b. Find an equation for  $C$  (4)

**(Total Marks: 6)**

---

8. Find, giving your answer to 3 significant figures where appropriate, the value of  $x$  for which,

a.  $3^x = 5$  (3)

b.  $\log_2(2x + 1) - \log_2 x = 2$  (4)

**(Total Marks: 7)**

---

9. Find all the values of  $x$ , to 1 decimal place, in the interval  $0^\circ \leq x < 360^\circ$  for which,

$5 \sin(x + 30) = 3$  (4)


b. Find all the values of  $x$ , to 1 decimal place, in the interval  $0^\circ \leq x < 360^\circ$  for which,

$\tan^2 x = 4$  (5)

**(Total Marks: 9)**

---

10. The figure shows the shaded region  $R$  which is bounded by the curve  $y = -2x^2 + 4x$  and the line  $y = \frac{3}{2}$ .  
The points  $A$  and  $B$  are the points of intersection of the line and the curve



Find,

a. The  $x$ -coordinates of the points  $A$  and  $B$  (4)

b. The exact area of  $R$ . (6)

**(Total Marks: 10)**

---

11.  $f(x) = 1 - \frac{3}{x+2} + \frac{3}{(x+2)^2}$ ,  $x \neq 2$

a. Show that  $f(x) = \frac{Ax^2 + Ax + A}{(x+2)^2}$ ,  $x \neq 2$ , where  $A$  is an integer to be found. (4)

b. Show that  $x^2 + x + 1 > 0$ , for all values of  $x$  (3)

c. Show that  $f(x) > 0$  for all values of  $x$ ,  $x \neq 2$  (1)

**(Total Marks: 8)**


---

12. The figure shows a sketch of part of the curve  $C$  with equation,

$$y = x(x - 1)(x - 5)$$



Use calculus to find the total area of the infinite region, shown shaded in the figure, that is between  $x = 0$  and  $x = 2$  and is bounded by  $C$ , the  $x$ -axis and the line  $x = 2$ . (9)



**(Total Marks: 9)**

13. A diesel lorry is driven from Birmingham to Bury at a steady speed of  $v$  kilometres per hour. The total cost of the journey,  $\pounds C$ , is given by

$$C = \frac{1400}{v} + \frac{2v}{7}$$

- a. Find the value of  $v$  for which  $C$  is a minimum (5)
- b. Find  $\frac{d^2C}{dv^2}$  and hence verify that  $C$  is a minimum for this value of  $v$  (2)
- c. Calculate the minimum total cost of the journey (2)

**(Total Marks: 9)**

14.  $f(x) = x^4 - x^3 + 3x^2 + ax + b$

Where  $a$  and  $b$  are constants.

When  $f(x)$  is divided by  $(x - 1)$  the remainder is 4

When  $f(x)$  is divided by  $(x + 2)$  the remainder is 22.

Find the value of  $a$  and the value of  $b$  (5)

**(Total Marks: 5)**

15. The radioactive decay of a substance is given by,

$$R = 1000e^{-ct}, t \geq 0$$

Where  $R$  is the number of atoms at time  $t$  years and  $c$  is a positive constant

- a. Find the number of atoms when the substance started to decay (1)
- It takes 5730 years for half of the substance to decay
- b. Find the value of  $c$  to 3 significant figures. (4)
  - c. Calculate the number of atoms that will be left when  $t = 22\,920$ . (2)

**(Total Marks: 7)**



16a. Prove that

$$\frac{\cos x}{1 + \sin x} + \frac{1 + \sin x}{\cos x} \equiv \frac{2}{\cos x} \quad (4)$$

b. Hence or otherwise, find for  $0 < x < 360$ , all the solutions of,

$$\frac{\cos x}{1 + \sin x} + \frac{1 + \sin x}{\cos x} = 4 \quad (2)$$

---

**(Total Marks: 6)**

---

**Total Marks: 120**

---



## Mark Scheme

<b>1a</b>	$y = 5 - (2 \times 3) = -1$	<b>B1</b>
<b>1b</b>	Gradient of $L = \frac{1}{2}$	<b>B1</b>
	$y - (-1) = \frac{1}{2}(x - 3)$	<b>M1</b> <b>A1</b>
	$x - 2y - 5 = 0$	<b>A1</b>

<b>2a</b>	$3\sqrt{5}$	<b>B1</b>
<b>2b</b>	$\frac{2(3+\sqrt{5})}{(3-\sqrt{5})} \times \frac{(3+\sqrt{5})}{(3+\sqrt{5})}$	<b>M1</b>
	$\frac{2(14+6\sqrt{5})}{4}$	<b>B1</b> <b>M1</b> <b>M1</b>
	$= 7 + 3\sqrt{5}$	<b>a1</b>

<b>3a</b>	Shape of $f(x)$	<b>B1</b>
	Moved up	<b>M1</b>
	Asymptotes $y = 3$	<b>B1</b>
	$x = 0$	<b>B1</b>
<b>3b</b>	$\frac{1}{x} + 3 = 0$	<b>M1</b>
	$x = -\frac{1}{3}$	<b>A1</b>

<b>4</b>	Use of $b^2 - 4ac$	<b>M1</b>
	$(-3)^2 - 4 \times 2 \times x - (k + 1) < 0$	<b>A1</b>
	$8k < -17$	<b>M1</b>
	$k < -\frac{17}{8}$	<b>A1</b>

<b>5a</b>	$4x \rightarrow kx^2$	<b>M1</b>
	$f(x) = 2x^2 - 4x^{\frac{3}{2}} - 8x^{-1} + C$	<b>A1</b> <b>A1</b> <b>A1</b>
	At $x = 4, y = 1$	
	$1 = (2 \times 16) - (4 \times 4^{\frac{3}{2}}) - (8 \times 4^{-1}) + c$	<b>M1</b>
	$c = 3$	<b>A1</b>
<b>5b</b>	$f'(4) = 16 - (6 \times 2) + \frac{8}{16} = \frac{9}{2}$	<b>M1</b>
	Gradient of normal is $-\frac{2}{9} = -\frac{1}{m}$	<b>M1</b>
	Equation of normal: $y - 1 = -\frac{2}{9}(x - 4)$	<b>M1</b>
	$2x + 9y - 17 = 0$	<b>A1</b>

<b>6a</b>	$(x + 2k)^2$	<b>M1</b>
	$(x + 2k)^2 - 4k^2 + (3 + 11k)$	<b>M1</b> <b>A1</b>
<b>6b</b>	$4k^2 - 11k - 3 = 0$ $(4k + 1)(k - 3) = 0$	<b>M1</b>
	$k = -\frac{1}{4}$ $k = 3$	<b>A1</b>
	Using $b^2 - 4ac < 0$ $4k^2 - 11k - 3 < 0$	<b>M1</b>
	$-\frac{1}{4} < k < 3$	<b>A1</b>
<b>6c</b>	Shape	<b>B1</b>
	Minimum in correct quadrant, no touching $x$ -axis, not on the $y$ -axis, no other maximum or minimum.	<b>B1</b>
	$(0, 14)$ marked	<b>B1</b>

<b>7a</b>	$\left(\frac{5+13}{2}, \frac{-1+11}{2}\right) = (9, 5)$	<b>M1</b> <b>A1</b>
<b>7b</b>	$r^2 = (9 - 5)^2 + (5 - -1)^2 = 52$	<b>M1</b>
	Equation of circle,	<b>M1</b>
	$(x - 9)^2 + (y - 5)^2 = 52$	<b>A1</b> <b>A1</b>

<b>8a</b>	$\log 3^x = \log 5$	<b>M1</b>
	$x = \frac{\log 5}{\log 3}$	<b>A1</b>
	$x = 1.46$	<b>A1</b>
<b>8b</b>	$\log_2 \left(\frac{2x+1}{x}\right) = 2$	<b>M1</b>
	$\frac{2x+1}{x} = 2^2$	<b>M1</b>
	$2x + 1 = 4x$	<b>M1</b>
	$x = \frac{1}{2}$	<b>A1</b>

<b>9a</b>	$\sin(x + 30) = \frac{3}{5}$	<b>B1</b>
	$x + 30 = 36.9$	<b>B1</b>
	$x + 30 = 143.1$	<b>M1</b>
	$x = 6.9$	<b>A1</b>
	$x = 113.1$	
<b>9b</b>	$\tan x = \pm 2$	<b>B1</b>
	$\tan x = 2$	<b>B1</b> <b>M1</b>
	$x = 63.4$	
	$x = 243.4$	
	$\tan x = -2$	<b>M1</b>


	$x = 116.6^0$	
	$x = 296.6^0$	<b>M1</b>

<b>10a</b>	$\frac{3}{2} = -2x^2 + 4x$	<b>M1</b>
	$4x^2 - 8x + 3 = 0$	<b>A1</b>
	$(2x - 1)(2x - 3) = 0$	<b>M1</b>
	$x = \frac{1}{2}$ $x = \frac{3}{2}$	<b>A1</b>
<b>10b</b>	Area of R = $\int_{\frac{1}{2}}^{\frac{3}{2}} (-2x^2 + 4x) dx - \frac{3}{2}$	<b>B1</b>
	$\int (-2x^2 + 4x) dx = [-\frac{2}{3}x^3 + 2x^2]$	<b>M1</b> <b>A1</b>
	$\int_{\frac{1}{2}}^{\frac{3}{2}} (-2x^2 + 4x) dx = \left(-\frac{2}{3} \times \frac{3^3}{2^3} + 2 \times \frac{3^2}{2^2}\right) - \left(-\frac{2}{3} \times \frac{1}{2^3} + 2 \times \frac{1}{2^2}\right)$	<b>M1</b> <b>M1</b>
	$= \frac{11}{6}$ Area of R = $\frac{11}{6} - \frac{3}{2} = \frac{1}{3}$	<b>A1</b>

<b>11a</b>	$f(x) = \frac{(x+2)^2 - 3(x+2) + 3}{(x+2)^2}$	<b>M1</b> <b>A1</b> <b>A1</b>
	$= \frac{x^2 + 4x + 4 - 3x - 6 + 3}{(x+2)^2}$ $= \frac{x^2 + x + 1}{(x+2)^2}$	<b>A1</b>
	Where A = 1	
<b>11b</b>	$x^2 + x + 1 = (x + \frac{1}{2})^2 + \frac{3}{4}$	<b>M1</b> <b>A1</b>
	Any squared value will be positive, therefore, expression is positive.	<b>A1</b>
<b>11c</b>	$f(x) = \frac{(x + \frac{1}{2})^2 + \frac{3}{4}}{(x+2)^2}$ Numerator is positive and as $x \neq -2$ , $(x + 2)^2 > 0$ , therefore denominator is positive. Hence, $f(x) > 0$	<b>B1</b>

<b>12</b>	$y = x(x^2 - 6x + 5)$ $y = x^3 - 6x^2 + 5x$	<b>M1</b> <b>A1</b>
	$\int (x^3 - 6x^2 + 5x) dx = \frac{x^4}{4} - \frac{6x^3}{3} + \frac{5x^2}{2}$	<b>M1</b> <b>A1</b>
	$[\frac{x^4}{4} - \frac{6x^3}{3} + \frac{5x^2}{2}]_0^1 = (\frac{1}{4} - 2 + \frac{5}{2}) - 0 = \frac{3}{4}$	<b>M1</b>
	$[\frac{x^4}{4} - \frac{6x^3}{3} + \frac{5x^2}{2}]_1^2 = (4 - 16 + 10) - \frac{3}{4} = -\frac{11}{4}$	<b>M1</b> <b>A1</b>
	Total area = $\frac{3}{4} + \frac{11}{4} = \frac{7}{2}$	<b>M1</b> <b>A1</b>

<b>13a</b>	$\frac{dc}{dv} = -1400v^{-2} + \frac{2}{7}$	<b>M1</b> <b>A1</b>
	$-1400v^{-2} + \frac{2}{7} = 0$	<b>M1</b>
	$v^2 = 4900$	<b>M1</b>
	$v = 70$	<b>A1</b>
<b>13b</b>	$\frac{d^2c}{dv^2} = 2800v^{-3}$	<b>M1</b>
	$v = 70$ , $\frac{d^2c}{dv^2} > 0$	<b>A1</b>



	Therefore minimum	
<b>13c</b>	$v = 70$ $C = \frac{1400}{70} + \frac{2 \times 70}{7}$	<b>M1</b>
	$C = 40$	<b>A1</b>
<b>14</b>	$f(x) = x^4 - x^3 + 3x^2 + ax + b$ $f(1) = 4$ $f(2) = 22$	<b>M1</b>
	$3 + a + b = 4$	<b>A1</b>
	$16 + 8 + 12 - 2a + b = 22$	<b>A1</b>
	Solving simultaneously, $a = 5$ $b = -4$	<b>M1</b> <b>A1</b>
<b>15a</b>	1000	<b>M1</b>
<b>15b</b>	$1000e^{-5730c} = 500$	<b>M1</b>
	$e^{-5730c} = \frac{1}{2}$	<b>A1</b>
	$-5730c = \ln \frac{1}{2}$	<b>M1</b>
	$c = 0.000121$	<b>A1</b>
<b>15c</b>	$R = 1000e^{-2290c} = 62.5$	<b>M1</b> <b>A1</b>
<b>16a</b>	$\frac{\cos x}{1 + \sin x} + \frac{1 + \sin x}{\cos x} = \frac{\cos^2 x + (1 + \sin x)^2}{(1 + \sin x) \cos x}$	<b>M1</b>
	$\frac{\cos^2 x + 1 + 2 \sin x + \sin^2 x}{(1 + \sin x)(\cos x)}$	<b>A1</b>
	$= \frac{2(1 + \sin x)}{(1 + \sin x) \cos x}$	<b>M1</b>
	$= \frac{2}{\cos x}$	<b>A1</b>
<b>16b</b>	$\cos x = \frac{1}{2}$	<b>M1</b>
	$x = 60^\circ, 300^\circ$	<b>A1</b>



## Topic List

<b>Q1</b>	Equations of lines
<b>Q2</b>	Simplifying surds
<b>Q3</b>	Reciprocal graph sketching
<b>Q4</b>	Roots
<b>Q5</b>	Integration and normal
<b>Q6</b>	Completing the square and graph sketching
<b>Q7</b>	Circles
<b>Q8</b>	Solving logarithms
<b>Q9</b>	Solving trig. equations
<b>Q10</b>	Area under curves
<b>Q11</b>	Algebraic fractions
<b>Q12</b>	Area of a curve
<b>Q13</b>	Maxima and minima
<b>Q14</b>	Factor theorem
<b>Q15</b>	Exponential modelling
<b>Q16</b>	Trig proof and solving trig

