


AS-Level Unit Test: Integration



1. Find $\int 3x^2 + 4x^5 - 7 dx$, giving each term in its simplest form. (2)
2. Given that $y = 2x^3 + \frac{3}{x^2}$, $x \neq 0$ find,
 - a. $\frac{dy}{dx}$ (2)
 - b. $\int y dx$, simplifying each term (1)
3. A curve has an equation which satisfies $\frac{dy}{dx} = kx(2x - 1)$ for all values of x . The point $P(2, 7)$ lies on the curve and the gradient of the curve at P is 9.
 - a. Find the value of the constant k (2)
 - b. Find the equation of the curve (5)
4. The curve C has equation $y = f(x)$, $x > 0$, and $f'(x) = 4x - 6\sqrt{x} + \frac{8}{x^2}$. Given that the point $P(4, 1)$ lies on C .
 - a. Find $f(x)$ and simplify your answer (4)
 - b. Find an equation of the normal to C at the point $P(4, 1)$ (4)
5. Find the exact value of $\int_2^1 \frac{2x^3 - 6x^3}{x^2} dx$ (4)
6. Evaluate $\int_1^8 \frac{1}{\sqrt{x}} dx$, giving your answer in the form $a + b\sqrt{2}$, where a and b are integers (5)
7. The Figure shows a sketch of part of the curve with equation, $y = 3x - x^{\frac{3}{2}}$, $x \geq 0$.



The finite region S , bounded by the x -axis and the curve, is shown shaded in the figure.

- a. Find $\int 3x - x^{\frac{3}{2}} dx$ (3)
- b. Hence, find the area of S . (5)

Total marks: 37

Mark Scheme

1.

$\int 3x^2 + 4x^5 - 7 dx = \frac{3x^3}{3} + \frac{4x^5}{5} - 7x + c$	M1
$= x^3 + \frac{4}{5}x^5 - 7x + c$	M1

2a.

$y = 2x^3 + \frac{3}{x^2} = 2x^3 + 3x^{-2}$	M1
$\frac{dy}{dx} = 6x^2 - 6x^{-3}$	M1

2b.

$y = 2x^3 + \frac{3}{x^2} = 2x^3 + 3x^{-2}$	
$\int y dx = \frac{2}{4}x^3 + \frac{3x^{-1}}{-1} = \frac{1}{2}x^3 - 3x^{-1} + c$	M1

3a.

$\frac{dy}{dx} = kx(2x - 1)$ When $x = 2$, $\frac{dy}{dx} = 9$	M1
$9 = k(2)(3)$ $k = \frac{3}{2}$	M1

3b.

$y = \frac{3}{2} \int x(2x - 1) dx$ $= \frac{3}{2} \int 2x^2 - 2 dx$	M1
$= \frac{3}{2} \left[\frac{2x^3}{3} - \frac{x^2}{2} \right] + c$	M1
$= x^3 - \frac{3}{4}x^2 + c$	M1
When $x = 2$, $y = 7$ $7 = 5 + c$	M1
$c = 2$	M1
$y = x^3 - \frac{3}{4}x^2 + 2$	

4a.

$y = f(x) = \int f'(x) dx$ $\int (4x - 6\sqrt{x} + \frac{8}{x^2}) dx = \frac{4x^2}{2} - \frac{6x^{\frac{3}{2}}}{\frac{3}{2}} + \frac{8x^{-1}}{-1} + c$	M1
$y = 2x^2 - 4x^{\frac{3}{2}} - \frac{8}{x} + c$	M1
When $f(4) = 1$, $2(4)^2 - 4(4)^{1.5} - 2 + c = 1$	M1
$32 - 32 - 2 + c = 1$ $c = 3$	M1
$y = 2x^2 - 4x^{\frac{3}{2}} - \frac{8}{x} + 3$	



4b.

$f'(x) = 4x - 6\sqrt{x} + \frac{8}{x^2}$ $x = 4,$ $f'(x) = \frac{9}{2}$	M1
Gradient of normal at P = $-\frac{2}{9}$	M1
Equation of normal at P is: $y - 1 = -\frac{2}{9}(x - 4)$	M1
$9y - 9 = -2(x - 4)$ $2x + 9y - 17 = 0$	M1

5.

$\frac{2x^3 - 6x^3}{x^2} = \frac{2x^3}{x^2} - \frac{6x^3}{x^2} = 2x - 6$	M1
$\int_2^1 \frac{2x^3 - 6x^3}{x^2} dx = \int_2^1 2x - 6 dx$ $= \left[\frac{2x^2}{2} - 6x \right]_2^1$ $= [x^2 - 6x]_2^1$	M1
$((2)^2 - 6(2)) - ((1)^2 - 6(1))$	M1
$= -5 - (-8)$ $= 3$	M1

6.

$\int_1^8 \frac{1}{\sqrt{x}} dx = \int_1^8 x^{-\frac{1}{2}} dx$	M1
$= \left[\frac{x^{\frac{1}{2}}}{\frac{1}{2}} \right]_1^8 = [2x^{\frac{1}{2}}]_1^8$	M1
$= 2(8)^{\frac{1}{2}} - 2(1)^{\frac{1}{2}}$	M1
$2\sqrt{8} - 2\sqrt{1}$ $= 2\sqrt{8} - 2$ $= 2(2\sqrt{2}) - 2$ $= 4\sqrt{2} - 2$	M1
In the correct form: $-2 + 4\sqrt{2}$ $a = -2$ $b = 4$	M1

7a.

$\int 3x - x^{\frac{3}{2}} dx = \frac{3x^2}{2} - \frac{x^{\frac{5}{2}}}{\frac{5}{2}} + c$	M1 M1 M1
---	---------------------------

7b.

$3x - x^{\frac{3}{2}} = 0$ $x(3 - x^{\frac{1}{2}}) = 0$	M1
$x = 0$ $3 - x^{\frac{1}{2}} = 0 \rightarrow x^{\frac{1}{2}} = 3 \rightarrow x = 9$	M1
$\left[\frac{3x^2}{2} - \frac{x^{\frac{5}{2}}}{\frac{5}{2}} \right]_0^9$	M1

$= \left(\frac{3(9)^2}{2} - \frac{(9)^{\frac{5}{2}}}{\frac{5}{2}} \right) - \left(\frac{3(0)^2}{2} - \frac{(0)^{\frac{5}{2}}}{\frac{5}{2}} \right)$	M1
$= \frac{243}{2} - \frac{486}{5}$ $= \frac{243}{10}$	M1

